

## **Simulation of the Elastin and Fibrillin in Non-Irradiated or UVA Radiated Fibroblasts, and Direct Inhibition of Elastase or Matrix Metalloproteinases Activity by Nicotinamide or Its Derivatives**

NEENA PHILIPS, JOVINNA CHALENSOUK-KHAOSAAT, and SALVADOR GONZALEZ, *Department of Biology, Fairleigh Dickinson University, Teaneck, NJ (N.P., J.C-K.) and Department of Dermatology, Medicine and Medical Specialties, Alcala University, Madrid, Spain (S.G.).*

*Accepted for publication November 28, 2017.*

### **Synopsis**

Skin aging/photoaging is associated with altered the structure of collagen and elastin fibers, and increased activity of matrix metalloproteinases (MMP) and elastase. Nicotinamide and its derivatives, 2,6-dihydroxynicotinamide, 2,4,5,6-tetrahydroxynicotinamide, and 3-hydroxypicolinamide (collectively niacin derivatives) stimulate fibrillar collagen and heat shock proteins in dermal fibroblasts. The goal of this research was to extend the understanding of the anti-skin aging mechanism of these niacin derivatives through the stimulation of elastin (at the protein and promoter levels), fibrillin (1 and 2) in nonirradiated or ultraviolet (UVA) radiated dermal fibroblasts, and through the direct inhibition of MMP (1, 3, and 9) and elastase activities. UVA radiation stimulated elastin and inhibited fibrillin-1 and fibrillin-2 in dermal fibroblasts. The niacin derivatives significantly stimulated the expression of elastin (transcriptionally), fibrillin-1 and fibrillin-2 in nonirradiated and UVA radiated fibroblasts, and directly inhibited MMP or elastase activity. Overall, the niacin derivatives, more so nicotinamide and 2,6-dihydroxynicotinamide, have anti-skin aging potential through the stimulation of elastin and fibrillin, and the direct inhibition of the extracellular matrix proteolytic enzymes.

### **INTRODUCTION**

The structural integrity of the extracellular matrix (ECM), composed predominantly of collagen and elastin fibers, is essential to skin structure and function (1–8). The structure of the ECM deteriorates with intrinsic aging, and exposure to environmental factors such as ultraviolet (UV) radiation (1–27). UVA radiation can penetrate the dermis and damage ECM through the generation of oxidative stress and inflammation, and direct damage of biomolecules (9–14). The dermal fibroblasts are the primary synthesizers of the ECM proteins (1–4,8).

---

Address all correspondence to Neena Philips at [nphilips@fdu.edu](mailto:nphilips@fdu.edu) and [neenaphilips@optonline.net](mailto:neenaphilips@optonline.net).

Polyphenols with their antioxidative and anti-inflammatory properties have been identified to beneficially regulate the ECM, and thereby, prevent skin aging or cancer (1–8,27). The structure of the phenolic components includes at least one aromatic ring with one or more hydroxyl groups (5). The activity of the phenolic compounds is dependent on the number and location of these hydroxyl groups, in addition to their structure (5,28–34). The structure of nicotinamide or its derivatives, such as 3-hydroxypicolinamide, has photophysical and UV radiation absorptive properties (28–34). Skin aging or photoaging reduces cellular antioxidant defense, ECM structure, and NAD content (1–8,35). Nicotinamide serves as a precursor for several cellular coenzymes that are essential to metabolism and counteracts inhibition of ATP/glycolysis, oxidative DNA damage, oxidative stress, and inflammation (35–41). Nicotinamide improves skin appearance, independently or in combination with retinoids, by reducing hyperpigmentation and wrinkles (42–46). We recently reported an anti-skin aging mechanism of nicotinamide through the stimulation of fibrillar collagen and heat shock proteins in dermal fibroblasts (27).

The goal of this research was to determine the anti-skin aging mechanism of nicotinamide and three of its derivatives, 2,6-dihydroxynicotinamide, 2,4,5,6-tetrahydroxynicotinamide, and 3-hydroxypicolinamide (collectively niacin derivatives), through the beneficial regulation of elastin and fibrillin (1 and 2) in nonirradiated or UVA-radiated dermal fibroblasts, and direct inhibition of on matrix metalloproteinases (MMP) (1, 3, and 9) and elastase activities. The hypothesis of this research was that the niacin derivatives would stimulate expression of elastin and fibrillin, and directly inhibit ECM proteolytic activity.

## METHODS

### CELL CULTURE AND DOSING

Human neonatal dermal fibroblasts from two donors (Cascade Biologics, part of Thermofischer Scientific, Waltham, MA) were cultured in complete Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% heat inactivated fetal bovine serum, 1% penicillin/streptomycin (P/S), and 1% L-glutamine (Sigma, St. Louis, MO). The cells were rinsed with Hank's Balanced Salt solution (Sigma) and nonirradiated (control) or radiated with 2.5 J/cm<sup>2</sup> (minimal toxicity) of UVA radiation using a four-tube UVA lamp with irradiance of 1E–03 W/cm<sup>2</sup> for a sum of wavelengths from 320 to 400 nm (2,4,27). The nonirradiated or UVA-radiated cells were then not exposed (control or UVA-radiated control) or exposed to 0.01% (0.05 mM), 0.1% (0.5 mM), or 1% (5 mM) of each of the niacin derivatives in experimental media (DMEM containing 1% serum replacement and 1% P/S) for 24 h (2,4,27). Four independent experiments, in replicates of 3–4, were performed with dermal fibroblasts at passages 4–12. The cells were examined for cell viability by MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium, inner salt) assay (Promega, Madison, WI). The cell viability was not altered at these concentrations of UVA radiation or niacin derivatives.

### ELASTIN, AND FIBRILLIN (-1, -2) PROTEIN LEVELS

The elastin and fibrillin proteins levels in the media of nonirradiated or UVA radiated fibroblasts ± niacin derivatives were determined by using enzyme linked immunosorbant assay (ELISA) (Kirkgaard and Perry Laboratories Inc., Milford, MA; Elastin Products Co.)

(1–4,8,27). One hundred microliter aliquots of media or cells from each sample, or respective standards were added to independent wells of 96 well plates for 24 h. The wells were blocked with bovine serum albumin, and then incubated with respective antibodies (Elastin Products Co.) for 1 h. The plates were washed with wash buffer, incubated with respective secondary antibodies linked to peroxidase for 1 h, washed, and subsequently incubated with peroxidase substrate until color development, which was measured spectrophotometrically at 405 nm and quantitated from standard curves.

#### ELASTIN PROMOTER ACTIVITY

Fibroblasts were cotransfected with elastin promoter-firefly luciferase (Pgl4 vector) and TK-Renilla luciferase plasmids (for normalization of transfection efficiency) using Escort (Sigma) for 24 h before dosing with or without UVA-radiation  $\pm$  niacin derivatives for 24 h (2,4,8). The cells were measured for luminescence from firefly or renilla luciferase with specific substrates and quantitated using recombinant luciferase as standard (Promega).

#### MMP-1, -9, AND -9, AND ELASTASE ACTIVITIES

The inhibition of ECM proteolytic enzymes (MMP-1, MMP-2, MMP-3, Elastase) (Biomol, Torrance, CA; Enzo Life Sciences, Farmingdale, NY; Elastin products Co.) was performed as previously reported (1,3). Each of the enzymes at optimal concentration was incubated with the niacin derivatives at 0%, 0.01%, 0.1%, or 1% for 10 min followed by the addition of its respective substrate (Bachem). The reaction kinetics were measured fluorometrically (355 excitation/450 emission) every 10 min for a total of 60 min. The initial reading (0 time) was subtracted from the final reading (60 min) and the data converted to percentage of control.

#### DATA ANALYSIS

The data were analyzed for significant difference by analysis of variance and student *t*-tests at 95% confidence interval. The effects of UVA radiation on dermal fibroblasts were analyzed relative to nonirradiated control cells. The effects of the niacin derivatives on nonirradiated cells were analyzed relative to nonirradiated cells (control). The effects of each of the niacin derivatives on UVA-radiated fibroblasts were analyzed relative to UVA radiation effect alone (UVA-radiated respective control). The direct MMP or elastase inhibitory activity of the niacin derivatives was analyzed relative to control.

## RESULTS

#### STIMULATION OF EXPRESSION OF ELASTIN, ELASTIN PROMOTER, FIBRILLIN-1, AND FIBRILLIN-2 BY NICOTINAMIDE, 2,6-DIHYDROXYNICOTINAMIDE, 2,4,5,6-TETRAHYDROXYNICOTINAMIDE, AND 3-HYDROXYPICOLINAMIDE IN NONIRRADIATED FIBROBLASTS

Niacin or its derivatives stimulated the expression of elastin and fibrillin in dermal fibroblasts. The niacin derivatives significantly stimulated elastin expression (protein and promoter) and fibrillin-1 and fibrillin-2 at 0.1% and 1% in nonirradiated fibroblasts ( $p < 0.05$ ,

relative to control) (Figure 1A–D). In addition, fibrillin-1 and fibrillin-2 were stimulated by nicotinamide and 2,6-dihydroxynicotinamide at 0.01% in nonirradiated fibroblasts ( $p < 0.05$ , relative to control) (Figure 1C and D).

In nonirradiated fibroblasts, the elastin protein levels (70 ng/ml as 100%)/elastin promoter activity (400 pg/ml as 100%) were stimulated by nicotinamide, 2,6-dihydroxynicotinamide, 2,4,5,6-tetrahydroxynicotinamide, and 3-hydroxypicolinamide upto 308/222%, 413/719%, 243/358%, and 224/183% of nonirradiated controls, respectively ( $p < 0.05$ ) (Figure 1A and B). The expression of fibrillin-1 (0.5  $\mu\text{g/ml}$  as 100%)/fibrillin-2 (35 ng/ml as 100%) was stimulated by nicotinamide, 2,6-dihydroxynicotinamide, 2,4,5,6-tetrahydroxynicotinamide, and 3-hydroxypicolinamide upto 614/578%, 601/533%, 402/234%, and 336/290% of nonirradiated controls, respectively ( $p < 0.05$ ) (Figure 1C and D).

STIMULATION OF EXPRESSION OF ELASTIN, ELASTIN PROMOTER, FIBRILLIN-1, AND FIBRILLIN-2 BY NICOTINAMIDE, 2,6-DIHYDROXYNICOTINAMIDE, 2,4,5,6-TETRAHYDROXYNICOTINAMIDE, AND 3-HYDROXYPICOLINAMIDE IN UVA RADIATED FIBROBLASTS

UVA-radiation stimulated elastin protein level to 185% ( $\pm 14\%$ ) of control, and elastin promoter activity to 125% ( $\pm 8\%$ ) of control ( $p < 0.05$ ). UVA-radiation significantly inhibited fibrillin-1 and fibrillin-2 to 65% ( $\pm 4\%$ ) of control and 73% ( $\pm 7\%$ ) of control ( $p < 0.05$ ). The effects of the niacin derivatives on UVA-radiated fibroblasts were similar




Figure 1. Stimulation of elastin protein (A), elastin promoter activity (B), fibrillin-1 protein (C), and fibrillin-2 protein (D) by nicotinamide (green line), 2,6-dihydroxynicotinamide (red line), 2,4,5,6-tetrahydroxynicotinamide (violet line), and 3-hydroxypicolinamide (blue line) in non-irradiated dermal fibroblasts; \* =  $p < 0.05$ , relative to control, error bars (A–D) represent standard deviation,  $n = 4$ .

to those on non-irradiated fibroblasts, except for the lack of regulation of fibrillin expression by 0.01% nicotinamide and 2,6-dihydroxynicotinamide.

Relative to UVA-irradiated controls, the expression of elastin protein/promoter was stimulated by nicotinamide, 2,6-dihydroxynicotinamide, 2,4,5,6-tetrahydroxynicotinamide, and 3-hydroxypicolinamide upto 348/268%, 365/708%, 230/547%, and 221/214%, respectively ( $p < 0.05$ ) (Figure 2A and B). The expression of fibrillin-1/fibrillin-2 protein levels was stimulated by nicotinamide, 2,6-dihydroxynicotinamide, 2,4,5,6-tetrahydroxynicotinamide, and 3-hydroxypicolinamide upto 524/482%, 859/563%, 347/312%, and 253/227% of UVA-irradiated controls, respectively ( $p < 0.05$ ) (Figure 2C and D).

DIRECT INHIBITION OF MMP-1, MMP-3, MMP-9, AND ELASTASE ACTIVITIES BY NICOTINAMIDE, 2,6-DIHYDROXYNICOTINAMIDE, 2,4,5,6-TETRAHYDROXYNICOTINAMIDE, AND 3-HYDROXYPICOLINAMIDE

The MMP-1 and MMP-3 activities were significantly inhibited by nicotinamide and 2,6-dihydroxynicotinamide; MMP-9 activity by nicotinamide, and elastase activity by nicotinamide, 2,6-dihydroxynicotinamide, 2,4,5,6-tetrahydroxynicotinamide, and 3-hydroxypicolinamide ( $p < 0.05$ ) (Figure 3).

The MMP-1/MMP-3 activities were significantly inhibited by nicotinamide upto 43/36%, and by 2,6-dihydroxynicotinamide upto 70/57%, of respective controls ( $p < 0.05$ )




Figure 2. Stimulation of elastin protein (A), elastin promoter activity (B), fibrillin-1 protein (C), and fibrillin-2 protein (D) by nicotinamide (green line), 2,6-dihydroxynicotinamide (red line), 2,4,5,6-tetrahydroxynicotinamide (violet line), and 3-hydroxypicolinamide (blue line) in UVA-irradiated dermal fibroblasts; \* =  $p < 0.05$ , relative to UVA irradiated control cells, error bars (A–D) represent standard deviation,  $n = 4$ .

(Figure 3A). There was direct stimulation of MMP-3 activity by 3-hydroxypicolinamide upto 329% of control, respectively ( $p < 0.05$ ) (Figure 3B).

The MMP-9 activity was significantly inhibited only by nicotinamide, upto 44% of control ( $p < 0.05$ ) (Figure 3C).

The elastase activity was inhibited by nicotinamide, 2,6-dihydroxynicotinamide, 2,4,5,6-tetrahydroxynicotinamide and 3-hydroxypicolinamide at 1–40%, 72%, 70%, and 66% of control, respectively ( $p < 0.05$ ) (Figure 3D).

## DISCUSSION

Nicotinamide, with an amide linked to an aromatic ring has UV absorptive, anti-inflammatory properties, cellular metabolic, and antiapoptotic properties. The hypothesis of this research was that nicotinamide and its derivatives, 2,6-dihydroxynicotinamide, 2,4,5,6-tetrahydroxynicotinamide, and 3-hydroxypicolinamide would stimulate elastin and fibrillin in nonirradiated, and UVA radiated dermal fibroblasts, and exhibit direct antiproteolytic activity on the ECM proteins.

The elastin fibers, composed of elastin and fibrillin microfibrils, give skin its firmness and elasticity (1–4,15–18). The formation of the elastin fibers in the dermis is directed by the microfibrils, composed primarily of fibrillin-1 and fibrillin-2. The microfibrils also form at the epidermal–dermal junction (15). There is loss of elastin and fibrillin with intrinsic aging, and in addition elastotic deposits occur in photoaged skin (1–4,15–18). Nicotinamide




Figure 3. Direct regulation of MMP-1 activity (A), MMP-3 activity (B), MMP-9 activity (C), and elastase activity (D) by nicotinamide (green line), 2,6-dihydroxynicotinamide (red line), 2,4,5,6-tetrahydroxynicotinamide (violet line), and 3-hydroxypicolinamide (blue line); \* =  $p < 0.05$ , relative to control, error bars (A–D) represent standard deviation,  $n = 4$ .

(4–5%), independently or in combination with peptide/retinyl propionate, reduces skin wrinkles and increases skin elasticity in subjects; and is better tolerated than tretinoin (42–44). The niacin derivatives stimulated the expression of elastin and fibrillin in nonirradiated and UVA radiated fibroblasts, with nicotinamide and 2,6-dihydroxynicotinamide having a more potent effect. It is inferred that niacin or its derivatives improve skin elasticity by stimulating the expression of elastin and fibrillin in dermal fibroblasts.

There is loss of elastin and fibrillin with intrinsic aging, and in addition elastotic deposits occur in photoaged skin (1–4,15–18). In dermal fibroblasts, UVA radiation stimulates elastin expression whereas UVB-radiation inhibits it (2). Topical 5% nicotinamide counteracts UV-radiation-induced immunosuppression in human subjects (37). Nicotinamide at 50  $\mu\text{m}$  prevents UV-radiation-induced oxidative damage in HaCat keratinocytes (40). UVA radiation stimulated the expression of elastin, and inhibited fibrillin-1 and fibrillin-2 protein levels in dermal fibroblasts. The expression of elastin, fibrillin-1, and fibrillin-2 were significantly stimulated by 0.1% and 1% each of the four niacin derivatives in UVA radiated fibroblasts, which suggests that the mechanism to nicotinamide's counteraction of the clinical signs of photoaging is through the stimulation and formation of proper elastin fibers. The increased expression of fibrillin-1 and fibrillin-2 may allow for deposition of well-formed elastic fibers instead of elastotic deposits.

The ECM is remodeled by MMPs and elastases (19–26). The proteolytic enzymes of collagen and elastin fibers include MMPs-1, -3, -9 and elastase (18–26). A mechanism to the damage to the ECM with skin aging is the increased activity of these ECM remodeling enzymes. The niacin derivatives inhibited elastase activity, and nicotinamide and 2,6-dihydroxynicotinamide inhibited the activities of MMPs, which suggests protective effects on the dermal collagen and elastin fibers.

## CONCLUSION

The clinical signs of intrinsic aging and photoaging result from increased oxidative stress, which activates mitogen activated protein kinase pathway and activator protein (AP)-1 transcription factor, and inflammation, which activates nuclear factor-kappa B (NF- $\kappa$ B) transcription factor (5–7,27). Nicotinamide reduces wrinkles and improves the appearance of skin (35–41). This research reports for the first time the anti-skin aging mechanism of nicotinamide and its derivatives; 2,6-dihydroxynicotinamide; 2,4,5,6-tetrahydroxynicotinamide; and 3-hydroxypicolinamide through the stimulation of elastin transcription and fibrillin expression in nonirradiated and UVA radiated fibroblasts; and the direct inhibition of ECM proteolytic enzymes. The transcription of elastin is inhibited by AP-1 and NF- $\kappa$ B transcription factors (47,48). It is inferred that the niacin derivatives reduce oxidative stress and inflammation, and thereby AP-1 and NF- $\kappa$ B transcription factors; to facilitate the increased transcription of the elastin gene. The formation of organized elastin fibers *in vivo* would need to be investigated to validate the ECM effect of nicotinamide in skin.

## ACKNOWLEDGMENTS

Marvin Tuason, Halyna Siomyk, and Aksinya Kogan contributed to the research through technical assistance, data analysis/presentation, or literature search. Funding was provided by Industrial Farmaceutica Cantabria (IFC), Spain.

## REFERENCES

- (1) N. Philips, M. Samuel, R. Arena, Y. Chen, J. Conte, P. Natrajan, G. Haas, and S. Gonzalez, Direct inhibition of elastase and matrixmetalloproteinases, and stimulation of biosynthesis of fibrillar collagens, elastin and fibrillins by xanthohumol, *J. Cosmet. Sci.*, **61**, 125–132 (2010).
- (2) N. Philips, J. Smith, T. Keller, and S. Gonzalez, Predominant effects of *Polypodium leucotomos* on membrane integrity, lipid peroxidation, and expression of elastin and matrixmetalloproteinase-1 in ultraviolet radiation exposed fibroblasts, and keratinocytes, *J. Dermatol. Res.*, **32**, 1–9 (2003).
- (3) N. Philips, P. Samuel, H. Parakandi, S. Gopal, H. Siomyk, A. Ministro, T. Thompson, and G. Borkow, Beneficial regulation of fibrillar collagens, heat shock protein-47, elastin fiber components, transforming growth factor- $\beta$ 1, vascular endothelial growth factor and oxidative stress effects by copper in dermal fibroblasts, *Connective Tissue Res.*, **53**, 373–378 (2012).
- (4) N. Philips, T. Keller, C. Hendrix, S. Hamilton, R. Arena, M. Tuason, and S. Gonzalez, Regulation of the extracellular matrix remodeling by lutein in dermal fibroblasts, melanoma cells, and ultraviolet radiation exposed fibroblasts, *Arch. Dermatol. Res.*, **299**, 373–379 (2007).
- (5) N. Philips, H. Siomyk, D. Bynum, and S. Gonzalez, “Skin Cancer, Polyphenols, and Oxidative Stress,” in *Cancer: Oxidative Stress and Dietary Antioxidants*. (Elsevier, London, U.K., 2013).
- (6) N. Philips, M. Samuel, H. Parakandi, H. Siomyk, S. Gopal, H. Jia, and H. Shahin, “Vitamins in the Therapy of Inflammatory and Oxidative Disease,” in *Frontiers in Clinical Drug Research-Anti Allergy Agents*. Bentham Science Publishers, Beijing, CN (2012).
- (7) N. Philips, P. Samuel, H. Siomyk, H. Parakandi, H. Jia, S. Gopal, and H. Shahin, “Improved Cell Metabolism and Strengthening of the Extracellular Matrix by Nicotinamide, and Copper for Anti-Skin Aging,” in *Skin Aging: New Research*, (Nova Science, New York, 2012).
- (8) N. Philips, J. Conte, Y. Chen, P. Natrajan, M. Taw, T. Keller, J. Givant, M. Tuason, L. Dulaj, D. Leonardi, and S. Gonzalez, Beneficial regulation of matrixmetalloproteinases and their inhibitors, fibrillar collagens and transforming growth factor-beta by *Polypodium leucotomos*, directly or in dermal fibroblasts, ultraviolet radiated fibroblasts, and melanoma cells, *Arch. Dermatol. Res.*, **301**, 487–495 (2009).
- (9) J. A. Nichols, and S. K. Katiyar, Skin photoprotection by natural polyphenols: Anti-inflammatory, antioxidant and DNA repair mechanisms, *Arch. Dermatol. Res.*, **302**, 71–83 (2010).
- (10) M. A. Bachelor, and G. T. Bowden, UVA-mediated activation of signaling pathways involved in skin tumor promotion and progression, *Semin. Cancer Biol.*, **14**, 131–138 (2004).
- (11) M. Gasparrini, T. Y. Forbes-Hernandez, S. Afrin, J. M. Alvarez-Suarez, A. M. González-Paramàs, C. Santos-Buelga, S. Bompadre, J. L. Quiles, B. Mezzetti, and F. Giampieri, A pilot study of the photoprotective effects of strawberry-based cosmetic formulations on human dermal fibroblasts, *Int. J. Mol. Sci.*, **16**, 17870–17884 (2015).
- (12) W. Min, X. Liu, Q. Qian, B. Lin, D. Wu, M. Wang, I. Ahmad, N. Yusuf, and D. Luo, Effects of baicalin against UVA-induced photoaging in skin fibroblasts, *Am. J. Chin. Med.*, **42**, 709–727 (2014).
- (13) C. Battie, S. Jitsukawa, F. Bernerd, S. Del Bino, C. Marionnet, and M. Verschoore, New insights in photoaging, UVA induced damage and skin types, *Exp. Dermatol.*, **23**, 7–12 (2014).
- (14) J. Ryu, S. J. Park, I. H. Kim, Y. H. Choi, and T. J. Nam, Protective effect of porphyra-334 on UVA-induced photoaging in human skin fibroblasts, *Int. J. Mol. Med.*, **34**, 796–803 (2014).
- (15) R. E. Watson, C. E. Griffiths, N. M. Craven, C. A. Shuttleworth, and C. M. KIELTY, Fibrillin-rich microfibrils are reduced in photoaged skin. Distribution at the dermal-epidermal junction, *J. Invest. Dermatol.*, **112**, 782–787 (1999).
- (16) R. E. Watson, N. M. Craven, S. Kang, C. J. Jones, C. M. KIELTY, and C. E. Griffiths, A short-term screening protocol, using fibrillin-1 as a reporter molecule, for photoaging repair agents, *J. Invest. Dermatol.*, **116**, 672–678 (2001).
- (17) M. J. Sherratt, Tissue elasticity and the ageing elastic fibre, *Age (Dordr.)*, **31**, 305–325 (2009).
- (18) N. Philips, “Experimental Physiology in Anti-Skin Aging,” in *Skin Anatomy and Physiology Research Development*, F. Columbus. Ed. (Nova Science Publishers Inc., New York, 2009).
- (19) M. R. Khorramizadeh, E. E. Tredget, C. Telasky, Q. Shen, and A. Ghahary, Aging differentially modulates the expression of collagen and collagenase in dermal fibroblasts, *Mol. Cell. Biochem.*, **194**, 99–108 (1999).
- (20) T. Quan, Z. Qin, W. Xia, Y. Shao, J. J. Voorhees, and G. J. Fisher, Matrix-degrading metalloproteinases in photoaging, *J. Invest. Dermatol. Symp. Proc.*, **14**, 20–24 (2009).
- (21) B. G. Godeau, H. Emonard, K. Baba, P. Bellon, W. Hornebeck, and G. Bellon, Analysis of the *ex vivo* specificity of human gelatinases A and B towards skin collagen and elastic fibers by computerized morphometry, *Matrix Biol.*, **19**, 139–148 (2009).



- (22) M. Brennan, H. Bhatti, K. C. Nerusu, N. Bhagavathula, S. Kang, G. J. Fisher, J. Varani, and J. J. Voorhees, Matrix metalloproteinase-1 is the major collagenolytic enzyme responsible for collagen damage in UV-irradiated human skin, *Photochem. Photobiol.*, **78**, 43–48 (2003).
- (23) F. Rijken, R. C. Kiekens, and P. L. Bruijnzeek, Skin-infiltrating neutrophils following exposure to solar-simulated radiation could play an important role in photoageing of human skin, *Br. J. Dermatol.*, **152**, 321–328 (2005).
- (24) J. Labat-Robert, A. Fourtanier, B. Boyer-Lafargue, and L. Robert, Age dependent increase of elastase type protease activity in mouse skin. Effect of UV-irradiation, *J. Photochem. Photobiol.*, **57**, 113–118 (2000).
- (25) K. Sellheyer, Pathogenesis of solar elastosis: Synthesis or degradation?, *J. Cutan. Patbol.*, **30**, 123–127 (2003).
- (26) T. M. Callaghan, and K. P. Wilhelm, A review of ageing and an examination of clinical methods in the assessment of ageing skin. Part 2: Clinical perspectives and clinical methods in the evaluation of ageing skin, *Int. J. Cosmet. Sci.*, **30**, 323–332 (2008).
- (27) N. Philips, J. Chalensouk-Khaosaat, and S. González, Stimulation of the fibrillar collagen and heat shock proteins by nicotinamide or its derivatives in non-irradiated or UVA radiated fibroblasts, and direct anti-oxidant activity of nicotinamide derivatives, *Cosmetics*, **2**, 146–161 (2015).
- (28) L. X. Cheng, J. J. Tang, H. Luo, X. L. Jin, F. Dai, J. Yang, Y. P. Qian, X. Z. Li, and B. Zhou, Antioxidant and antiproliferative activities of hydroxyl-substituted Schiff bases, *Bioorg. Med. Chem. Lett.*, **20**, 2417–2420 (2010).
- (29) Z. Y. Chen, P. T. Chan, K. Y. Ho, K. P. Fung, and J. Wang, Antioxidant activity of natural flavonoids is governed by number and location of their aromatic hydroxyl groups, *Chem. Phys. Lipids*, **79**, 157–163 (1996).
- (30) R. Gazák, P. Sedmera, M. Vrbacký, J. Vostálová, Z. Drahota, P. Marhol, D. Walterová, and V. Kren, Molecular mechanisms of silybin and 2,3-dehydrosilybin antiradical activity—role of individual hydroxyl groups, *Free Radic. Biol. Med.*, **46**, 745–758 (2009).
- (31) I. L. Rosa, P. C. Filho, C. R. Neri, O. A. Serra, A. T. de Figueiredo, J. A. Varela, and E. Longo, Synthesis and study of the photophysical properties of a new Eu<sup>3+</sup> complex with 3-hydroxypicolinamide, *J. Fluoresc.*, **21**, 1575–1583 (2011).
- (32) K. Sakai, T. Imakubo, M. Ichikawa, and Y. Taniguchi, A photoluminescent six-coordinated zinc(II) complex with hydroxides as axial ligands, [Zn(Hhpa)2(OH)2] (Hhpa = 3-hydroxypicolinamide), *Dalton Trans.*, **21**, 881–883 (2006).
- (33) M. Largeron and M. B. Fleury, Acid-base properties of pristinamycin IA and related compounds, *J. Pharm. Sci.*, **81**, 565–568 (1992).
- (34) Q. Wessels, E. Pretorius, C. M. Smith, and H. Nel, The potential of a niacinamide dominated cosmeceutical formulation on fibroblast activity and wound healing *in vitro*, *Int. Wound J.*, **11**, 152–158 (2014).
- (35) J. E. Oblong, The evolving role of the NAD<sup>+</sup>/nicotinamide metabolome in skin homeostasis, cellular bioenergetics, and aging, *DNA Repair (Amst.)*, **23**, 59–63 (2014).
- (36) P. A. Grange, J. Raingeaud, V. Calvez, and N. Dupin, Nicotinamide inhibits *Propionibacterium acnes*-induced IL-8 production in keratinocytes through the NF-kappaB and MAPK pathways, *J. Dermatol. Sci.*, **56**, 106–112 (2009).
- (37) D. L. Damian, C. R. S. Patterson, M. Stapelberg, J. Park, R. S. Barnetson, and G. M. Halliday, Ultraviolet radiation-induced immunosuppression is greater in men and prevented by topical nicotinamide, *J. Invest. Dermatol.*, **128**, 447–454 (2008).
- (38) K. Maiese, Z. Z. Chong, J. Hou, and Y. C. Shang, The vitamin nicotinamide: Translating nutrition into clinical care, *Molecules*, **14**, 3446–3485 (2009).
- (39) J. Park, G. M. Halliday, D. Surjana, and D. L. Damian, Nicotinamide prevents ultraviolet radiation-induced cellular energy loss, *Photochem. Photobiol.*, **86**, 942–948 (2010).
- (40) B. C. Thompson, G. M. Halliday, and D. L. Damian, Nicotinamide enhances repair of arsenic and ultraviolet radiation-induced DNA damage in HaCaT keratinocytes and *ex vivo* human skin, *PLoS One*, **10**, e0117491 (2015).
- (41) K. S. Ahn, K. Y. Moon, J. Lee, and Y. S. Kim, Downregulation of NF-kappaB activation in human keratinocytes by melanogenic inhibitors, *J. Dermatol. Sci.*, **31**, 193–201 (2003).
- (42) D. L. Bissett, J. E. Oblong, and C. A. Berge, Niacinamide: A B vitamin that improves aging facial skin appearance, *Dermatol. Surg.*, **31**, 860–865 (2005).
- (43) J. J. Fu, G. G. Hillebrand, P. Raleigh, J. Li, M. J. Marmor, V. Bertucci, P. E. Grimes, S. H. Mandy, M. I. Perez, S. H. Weinkle, and J. R. Kaczvinsky, A randomized, controlled comparative study of the

- wrinkle reduction benefits of a cosmetic niacinamide/peptide/retinyl propionate product regimen vs. a prescription 0.02% tretinoin product regimen, *Br. J. Dermatol.*, **162**, 647–654 (2010).
- (44) A. Kawada, N. Konishi, N. Oiso, S. Kawara, and A. Date, Evaluation of anti-wrinkle effects of a novel cosmetic containing niacinamide, *J. Dermatol.*, **35**, 637–642 (2008).
- (45) A. Pinkas-Sarafova, N. G. Markova, and M. Simon, Dynamic changes in nicotinamide pyridine dinucleotide content in normal human epidermal keratinocytes and their effect on retinoic acid biosynthesis, *Biochem. Biophys. Res. Commun.*, **336**, 554–564 (2005).
- (46) M. R. Namazi, Nicotinamide-containing sunscreens for use in Australasian countries and cancer-provoking conditions, *Med. Hypotheses*, **60**, 544–545 (2003).
- (47) V-M. Kahari, Y. Q. Chen, M. M. Bashir, J. Rosenbloom, and J. Uitto, Tumor necrosis factor-alpha down-regulates elastin gene expression. Evidence for the role of AP-1 in the suppression of promoter activity, *J. Biol. Chem.*, **267**, 26134–26141 (1992).
- (48) P. P. Kung, J. L. Berk, D. C. Rishikof, J. A. Foster, D. E. Humphries, D. A. Ricupero, R. H. Goldstein, NF-kappaB induced by IL-1beta inhibits elastin transcription and myofibroblast phenotype, *Am. J. Physiol. Cell Physiol.*, **283**, C58–C65 (2002).